

OR-10. RING-OPENING OF AZIRIDINES IN PRESENCE OF DIFFERENT REAGENTS AND CATALYSTS

A. Majee¹, S. Santra², S. Mahato¹, R. Chatterjee¹, G. V. Zyryanov^{2,3}

¹Department of Chemistry, Visva-Bharati (A Central University), Santiniketan, 731235, India

²Ural Federal University of the first President of Russia B. N. Yeltsin,
Mira St., 19, Yekaterinburg, 620002, Russia

³I. Ya. Postovsky Institute of Organic Synthesis UB RAS,
S. Kovalevskoy/Akademicheskaya St., 20/22, Yekaterinburg, 620990, Russia

E-mail: adinath.majee@visva-bharati.ac.in

Aziridines are highly versatile intermediates in organic synthesis due to their easy access and their susceptibility to ring-opening by facile C–N bond cleavage [1]. These ring opening products are 1,2-bifunctional compounds like diamines, aminols, amino ether, melatonin and α -amino-acid like tryptophan derivatives. In this lecture, it will be discussed our findings as an efficient and facile regioselective ring-opening of aziridines by various nucleophiles under mild and solvent-free conditions using zwitterionic-type salt [2]. We have also observed a regio-selective nucleophilic ring opening of aziridines by HCOOH where the HCOO[−] ion acts as a nucleophile [3]. We found that combination of NH₂OH · HCl and NaIO₄ is also very effective reagents for nucleophilic ring opening of aziridines [4].

References

1. Coleman R. S., Chen W. A Convergent Approach to the Mitomycin Ring System // Org. Lett. 2001. Vol. 3, № 8. P. 1141–1144.
2. Organocatalysis by an aprotic imidazolium zwitterion: regioselective ring-opening of aziridines and applicable to gram scale synthesis / Ghosal N. Chakraborty [et al.] // Green Chem. 2016. Vol. 18, № 2. P. 565–574.
3. An Efficient Synthesis of Oxazolidines by Tandem Ring-Opening / Closing Reaction of Ts-Aziridine Using Formic Acid / Ghosal N. Chakraborty [et al.] // Chemistry Select. 2018. Vol. 3, № 37. P. 10509–10514.
4. A Mild and Efficient Method for the Syntheses and Regioselective Ring-Opening of Aziridines / Ghosal N. Chakraborty [et al.] // SynOpen. 2017. Vol. 1, № 1. P. 0015–0023.

A. Majee acknowledges the financial support from DST-RSF Major Research Project (Ref. № INT/RUS/RSF/P-08). This work was supported by the Russian Science Foundation (Ref. № 18-73-00301).